

## Normal peak nasal inspiratory flow rate values in Greek children and adolescents

Papachristou A, Bourli E, Aivazi D, Futzila E, Papastavrou Th, Konstandinidis Th, Maratou E, Ilonidis G, Aivazis V

First Propedeutic Department of Internal Medicine, AHEPA Hospital, Aristotle University of Thessaloniki, Greece

### Abstract

**Background and aim:** Peak Nasal Inspiratory Flow Rate (PNIFR) is a clinical trial that has been instituted in clinical practice in order to determine the extent of nasal airway patency and it is used to assess the degree of nasal obstruction. This study attempts to provide tables referring to normal values of PNIFR in children and adolescents.

**Patients and Methods:** Three thousand one hundred and seventy pupils aged between 5 – 18 years, were selected to enter the study. Children with acute or chronic upper airway obstruction, such as acute obstructive pulmonary disease or allergic rhinitis and children below the 3<sup>rd</sup> percentile for weight and/or height were excluded from the study. All children that took part in the study were subjected to PNIFR measurements by using a portable Youtlen Peak Flow meter.

**Results:** A continuous increase of PNIFR values for boys and girls in relation to age increase was recorded. PNIFR values were higher in boys compared to girls and this difference was statistically significant until the age of 12.

**Conclusion:** Normal ranges for PNIFR standards are of great importance for the study of nasal patency, evaluation of the degree of nasal obstruction and application of treatment. This is the first time that a detailed description of PNIFR standards becomes available for the Greek population of children and adolescents. Hippokratia 2008; 12 (2): 94-102

**Key words:** nasal inspiratory flow rate, normal values, children, adolescents

**Corresponding author:** Aivazis V Th, 1<sup>st</sup> Department of Pediatrics, Aristotle University of Thessaloniki, 39 Diikitirou str., 546 30 Thessaloniki, Greece, Tel: 00302310-535240, Fax 00302310-533669, e-mail: aivazis@med.auth.gr

Nasal obstruction, may be related to age, body position, rhinic cycle or may be due to the presence of infection (tonsillitis, infection of adenoids), tonsillar or adenoidal hypertrophy, nasal polyps and allergies<sup>1</sup>. A positive family history should also be considered as an important risk factor of nasal obstruction<sup>2</sup>. Rhinoscopy is a reliable examination in order to decide the degree of nasal patency. The measurement of Peak Expiratory Flow Rate (PEFR) is a non-specific way to evaluate the presence of obstructive pulmonary disease and its response to treatment. The Peak Nasal Inspiratory Flow Rate (PNIFR) is an objective measurement for nasal obstruction and response to treatment regardless the etiology<sup>3-8</sup>.

It is a useful test for patients suffering from allergic rhinitis, for establishing the diagnosis and for monitoring treatment efficacy<sup>4,9</sup> and furthermore, it is a useful clinical tool to study various environmental factors that may cause nasal obstruction at home or at work<sup>10</sup>, or for deciding bronchial asthma treatment<sup>11</sup>.

In this retrospective study, we estimated the normal range of PNIFR values in Greek children according to age and sex and established the standards for Greek children.

### Patients – Methods

Four thousand one hundred eleven children (2010

boys and 2101 girls) aged between 5-18 years were examined. Children younger than 5 years old were excluded, as they could not follow the instructions. Of them 941 children (489 boys and 462 girls) or 22.88% (24.32% of the boys and 21.51% of the girls) were excluded from the study (Table 1). Eligibility was based on the European

**Table 1.** Needed criteria to include children and adolescents to the study according to European Respiratory Society (ERS).

1. Absence of acute disease.
2. No chest malformations, congenital abnormalities or respiratory tract diseases.
3. No cardiovascular or neuromuscular system diseases.
4. Free from symptoms of allergic rhinitis for the last 12 months.
5. Somatometric parameters above the 3<sup>rd</sup> percentile.

Respiratory Society (ERS) criteria for population studies<sup>12</sup> (Table 2).

Three thousand one hundred and seventy (3170) children (1521 boys and 1649 girls) were selected for the

**Table 2.** Number of children who took part, excluded from or included in the statistical analysis of our study according to sex.

	Number of children participating	Number of children excluded	Number of children included
<b>BOYS</b>	2010	489 (24.32 %)	1521
<b>GIRLS</b>	2101	452 (21.51 %)	1649
<b>TOTAL</b>	4111	941 (22.88 %)	3170

study. Authorities of the primary and high school supported the survey. Parental consensus was obtained. Parents were informed about the study and had to answer a questionnaire 2 or 3 days before their child's clinical examination. Following the recommendations of ISAAC (International Study of Allergy and Asthma in Children)<sup>13,14</sup> the questions were simply-worded and understandable, without the use of any medical terms. Each clinical examination took place in a classroom that was properly ventilated and had the appropriate temperature of 22-23°C. Children had to rest for at least 30 minutes before the examination.

The children were dressed with their usual clothes, did not wear shoes and were in upright position when subjected to the test, as suggested in current literature<sup>15</sup>.

The Seca Model 713 was used for height and weight measurement, as proposed by previous studies<sup>16</sup>.

PNIFR measurements were performed with a Youtlen Peak flow meter<sup>9</sup>, which is similar to a mini-Wright flow meter. Peak Nasal Inspiratory Flow Rate expressed in L/min is defined as the maximal instantaneous airflow achieved during forced inspiration through the nose. Asking the patient to take a deep, quick forced inspiration after having expired normally, performs the test. The amount of air left in the lungs after a tidal breath out is the Functional Residual Volume (FRV). Then the patient is instructed to inspire deeply through the nose, so that the Total Lung Capacity (TLC) is achieved. Total Lung Capacity (TLC) is the total volume of gas contained in the lungs at the end of a maximal inspiration<sup>1,3</sup>. The apparatus function was demonstrated and each pupil was instructed how to inhale forcefully. It is known that the number of attempts a child has to do in order to achieve the best score is inversely proportional to the child's age. Younger children may achieve the best score after the fourth or fifth attempt. The literature indicates that 5 repeated nasal inspirations do not obstruct nose vessels and do not affect the child's score. In our study, all children made 3 attempts of inspiration but only the best score was registered for each child as recommended in the literature<sup>9,17-23</sup>. The Youtlen Peak flow meter is a rather inexpensive portable device which is easy to use and which should be available, in clinical

practice, to all physicians.

Statistical analysis of recorded PNIFR included the estimation of mean, SD, SE and median values corresponding to the 50<sup>th</sup> percentile value of PNIFR. Curves for the 3<sup>rd</sup>, 5<sup>th</sup>, 95<sup>th</sup> and 97<sup>th</sup> percentiles values of PNIFR were calculated too<sup>24</sup>. Analysis of variance (ANOVA) was used to test hypothesis concerning means. Following that, correlation coefficients were calculated and the statistical significance of the Pearson correlation coefficient (R) was separately evaluated for boys, girls and for the total number of subjects.  $P < 0.05$  was considered to be significant.

## Results


The mean values of PNIFR, SD, SE and 95% confidence intervals for boys and girls according to their age are shown in Table 3.

PNIFR values for each percentile for boys and girls in relation to their age are shown in Table 4. A continuous increase of PNIFR values for boys and girls was observed in relation to age increase.

Figure 1 represents the mean PNIFR values for boys and girls when establishing 95% limits of significance. Boys achieved slightly greater scores compared to girls. The difference between boys and girls became statistically significant after the age of 12 ( $p < 0.05$ ).

**Table 3.** Number of children of each group (No), mean, standard deviation (sd), standard error (se) and the 95% confidence interval (CI) for PNIFR (L/min) for boys (B) and girls (G) according to their age (per year).

Age	Sex	No	Mean	Sd	Se	95%CI	
5-6	B	24	79.6	28.1	5.7	67.7	91.4
	G	23	80.4	29.9	6.2	67.5	93.4
6-7	B	99	100.1	34.2	3.4	93.3	106.9
	G	98	95.3	31.6	3.2	88.9	101.6
7-8	B	106	110.9	29.5	2.9	105.3	116.6
	G	128	100.0	30.9	2.7	96.6	108.5
8-9	B	105	115.0	32.8	3.2	108.7	121.3
	G	110	107.6	33.3	3.2	101.3	113.9
9-10	B	126	118.7	37.2	3.3	112.1	125.2
	G	120	110.4	38.7	3.5	103.4	117.4
10-11	B	117	131.4	35.3	3.3	124.9	137.9
	G	145	126.9	39.2	3.3	120.4	133.3
11-12	B	158	149.8	38.0	3.0	143.9	155.8
	G	136	145.0	40.9	3.5	138.1	152.0
12-13	B	148	168.7	27.4	2.3	164.3	173.2
	G	170	155.6	23.6	1.8	150.0	162.2
13-14	B	147	181.9	30.9	2.5	176.9	186.9
	G	179	160.2	29.2	2.2	155.9	165.5
14-15	B	171	190.0	33.9	2.6	184.9	195.1
	G	160	164.6	30.2	2.4	159.8	169.3
15-16	B	146	199.0	42.5	3.5	192.0	205.9
	G	180	168.7	33.4	2.5	162.8	173.6
16-17	B	125	204.3	40.3	3.6	197.2	211.4
	G	140	170.9	33.5	2.8	165.3	176.5
17-18	B	49	205.8	35.2	5.0	195.7	215.9
	G	60	172.5	30.4	3.9	164.6	179.4



**Figure 1.** Mean and 95% confidence interval (CI) for PNIFR according to the age (years) for both sexes (boys I. girls I).

**Discussion**

Result interpretation, following examinations of the respiratory system, requires the standards of the reference population. Reference of measured variables to population standards will help doctors to decide if there is aberration from the normal. Accurately defined population standards are of great importance.

The term standardized values has been widely used and has substituted the term normal values. The term standardized has been considered as correct since it is very difficult to define accurately the concept of being normal. Measurements characterized as standardized reflect the effort to establish new values and to modify old

tables, because the somatometric parameters of the individuals change as time goes by<sup>14,25</sup>.

Variable deviation from the normal may be attributed to many factors (technological, biological, environmental)<sup>26,27</sup>. Technological factors that may alter a result may be attributed to the accuracy of the apparatus, to the cooperation between the doctor and the patient and to the posture of the body and head’s position during the clinical examination<sup>26,28</sup>. Biological factors affecting the normal values are somatometric parameters such as weight, height and race<sup>25-29</sup>. Finally, environmental factors may also be considered, such as the region’s altitude, passive smoking and socioeconomic background<sup>30</sup>.

During the last few years many researchers resorted to the PNIFR for evaluating nasal patency by using Youlten Peak Flow meter and they have proved that this method is reliable<sup>4,9-11</sup>. Many recent studies have shown that the two techniques are similar, easy to perform and inexpensive<sup>18,19,31-33</sup>. According to Wihl et al<sup>31</sup> repeated PNIFR measurements had a difference of 5 L/min the one from another. Gleeson et al<sup>18</sup> similarly to rhinomanometry have also demonstrated high accuracy in PNIFR measurements.

In clinical practice these measurements are used in order to confirm diagnosis of a respiratory tract disease or to monitor treatment efficacy. PNIFR population measurements are considered normally distributed data and the sampling distribution represented in a normal curve can be used to test hypothesis about means. Confidence intervals for each sample mean is the sample mean plus or minus 2 times the standard deviation<sup>24,34</sup> for the 95% confidence interval (mean ± 2sd). So, 95% of the normally distributed cases lie within the confidence limits. When a value lies out this confidence interval, then percentiles should be used in order to characterize the result as normal or abnormal.

Our results have shown that PNIFR values tend to increase in proportion with age for boys and girls. There is no statistically significant difference in the mean value of one age group (annual distribution) compared to the next age group between children of the same sex (Graph 1). The increase is in a small range in girls after the age of 13 years, while in boys it keeps increasing in the same range until the age of 17-18 years according to our study. This may be attributed to the fact that girls do not grow taller after the age of 13 years. Other investigators, who found that lung volumes increase in relation to age, reported similar observation. Aivazis et al<sup>28</sup> and Barbarousis et al<sup>35</sup> reported that lung volumes in girls stop increasing after the age of 14 years and it was attributed to small changes of their height after this age<sup>25, 27, 28</sup>. Comparing PNIFR values between boys and girls we see that boys, in all age groups, have higher scores compared to girls, with the exception of the 5-6 years age group where PNIFR values for girls are slightly higher compared to those for boys.

**Table 4.** Number of children (No) of each group and percentiles for PNIFR (L/min) for boys and girls according to their age (per year).

Age	Sex	No	3 <sup>rd</sup> percentile	5 <sup>th</sup> percentile	50 <sup>th</sup> percentile	95 <sup>th</sup> percentile	97 <sup>th</sup> percentile
5-6	B	24	40.0	42.5	75.0	120.0	140.0
	G	23	40.0	50.0	80.0	130.0	135.0
6-7	B	99	50.0	55.0	95.0	140.0	150.0
	G	98	45.0	55.0	90.0	140.0	145.2
7-8	B	106	60.0	65.0	110.0	150.0	157.9
	G	128	50.0	60.5	95.0	150.0	155.0
8-9	B	105	65.9	70.0	115.0	160.0	170.0
	G	110	55.7	62.8	100.0	160.0	165.0
9-10	B	126	70.1	75.0	120.0	170.0	180.0
	G	120	60.0	65.0	110.0	170.5	175.0
10-11	B	117	75.0	80.5	140.0	185.0	195.0
	G	145	66.3	70.0	120.0	180.0	190.0
11-12	B	158	80.0	89.8	157.5	200.0	210.0
	G	136	70.0	80.0	150.0	200.0	205.9
12-13	B	148	110.0	124.5	170.0	220.0	230.0
	G	170	90.0	90.0	160.0	205.0	208.0
13-14	B	147	118.8	130.0	180.0	230.0	245.6
	G	179	95.0	100.0	170.0	210.0	220.0
14-15	B	171	125.0	140.0	190.0	250.0	270.0
	G	160	100.0	110.0	175.0	220.0	230.0
15-16	B	146	128.2	145.5	195.0	266.5	290.9
	G	180	102.0	115.0	180.0	230.0	235.7
16-17	B	125	135.0	150.0	200.0	277.0	302.2
	G	140	104.0	120.0	185.0	235.5	240.0
17-18	B	49	140.0	155.0	205.0	280.0	320.0
	G	60	104.9	120.5	190.0	239.5	243.1

PNIFR values for boys becomes statistically significantly higher after the age of 12 years ( $p < 0.001$ ).

To our knowledge, there are no references in Greek and international literature concerning PNIFR standards for children until now. All references available concern small groups of adults who had treatment with different drugs, especially those suffering from diseases that affect patency of the upper respiratory track system<sup>36-40</sup>. We, therefore, consider this study as the first internationally that attempts to assign PNIFR standards for children and adolescents in Greece and to provide an objective measure to all physicians for evaluating nasal airway patency in a relatively simple and inexpensive manner.

## References

- Pertuze J, Watson A, Pride NB. Maximum airflow through the nose in humans. *J Appl Physiol* 1991; 70:1369-1376
- Fairley JW, Durham LH, Ell SR. Correlation of subjective sensation of nasal patency with nasal inspiratory peak flow rate. *Clin Otolaryngol* 1993; 18:19-22
- Quanjer RhH, Tammeling GJ, Cotes JE, Pedersen OF, Peslin R. Lung volumes and forced ventilatory flows. *Pneumon* 1993; 6:11-43
- Youlten LJF. Nasal Airway Patency Measurement in the Assessment of Rhinitis Therapy. *Arbeiten aus dem Paul-Ehrlich-Institut, dem Georg-Speyer-Haus und dem Ferdinand-Blum-Institut. Heft 78 Gustav Fisher Verlag, Stuttgart. New York, 1983; pp. 111-113*
- Ahlstrom Emanuelson C, Anderson M, Persson CG, Thorsson L, Greiff L. Effects of topical formoterol alone and in combination with budesonide in a pollen season model of allergic rhinitis. *Respir Med* 2007 (E pub ahead of print)
- Can D, Tanac R, Demir E, Gulen F, Veral A. Is the usage of intranasal glucocorticosteroids alone in allergic rhinitis sufficient? *Allergy Asthma Proc* 2006; 27: 248-253
- Barnes MI, Ward JH, Fardon TC, Lipworth BJ. Effects of levocetirizine as add-on therapy to fluticasone in seasonal allergic rhinitis. *Clin Exp Allergy* 2006; 36:676-684
- Mim M, Sussman G, Hebert J, Lumry W, Lutsky B, Gates D. Desloratadine therapy for symptoms associated with perennial allergic rhinitis. *Ann Allergy Asthma Immunol* 2006; 96:460-465
- Youlten LJF. The peak nasal inspiratory flow meter: a new instrument for the assessment of the response to immunotherapy in seasonal allergic rhinitis. *Allergol Immunopathol* 1980; 8:344-347
- Nielsen J, Welinder H, Ottosson H, Bensryd I, Venge P, Skerfving S. Nasal challenge shows pathogenetic relevance of specific IgE serum antibodies for nasal symptoms caused by hexahydrophthalic anhydride. *Clin Exp Allergy* 1994; 24:440-449
- Pedersen S, Hansen OR, Fuglsang G. Influence of inspiratory flow rate upon the effect of a Turbuhaler. *Arch Dis Child* 1990; 65:308-310
- Quanjer RhH, Stocks J, Polgar J, Wise M, Karlberg J, Borsboom G. Complication of reference values for lung function measurements in children. *Eur Respir J* 1989; Suppl. 4:184s-261s
- Gratziou Ch, Priftis K, Anagnostakis I, et al. Prevalence of Asthma-Like Symptoms Among Greek Children. *Annual Congress of European Respiratory Society (ERS) 1996: 228s (abstract)*
- ISAAC. Steering Committee. Worldwide variations in prevalence of symptoms of asthma, allergic rhinoconjunctivitis and atopic eczema. *Lancet* 1998; 351:1225-1232
- Ogden C, Kuczmarski RJ, Flegal KM, et al. Centers for Disease Control and prevention 2000 growth charts for the United States: Improvements to the 1977 National Center for Health statistics version. *Pediatrics* 2002; 109:45-60
- Aivazis V. The normal child. Standards from 16.700 Greek children 0-17 years. *Monography. Thessaloniki 1990*
- Gardner R, Baker C, Broennle M, et al. American Thoracic Society (ATS). Statement-Snowbird workshop on standardization of spirometry. *Am Rev Respir Dis* 1979; 119:831-838
- Gleeson MJ, Youlten LJF, Shelton DM, Siodlak MZ, Eiser NM, Wengraf CL. Assessment of nasal airway patency: a comparison of four methods. *Clin Otolaryngol* 1986; 11:99-107
- Holmström M, Scadding GK, Lund VJ, Darby YC. Assessment of nasal obstruction. A comparison between rhinomanometry and nasal inspiratory peak flow. *Rhinology* 1990; 28:191-196
- Jones AS, Viani L, Phillips D, Charters P. The objective assessment of nasal patency. *Clin Otolaryngol* 1991; 16:206-211
- Clarke RW, Jones AS, Richardson H. Peak nasal inspiratory flow-the plateau effect. *J Laryngol Otol* 1995; 109:399-402
- Wilson AM, Dempsey OJ, Sims EJ, Lipworth BJ. Subjective and objective markers of treatment response in patients with seasonal allergic rhinitis. *Ann Allergy Asthma Immunol* 2000; 85:111-114
- Wilson AM, Dempsey OJ, Sims EJ, Coutie WJR, Paterson MC, Lipworth BJ. Evaluation of treatment response in patients with seasonal allergic rhinitis using domiciliary nasal peak inspiratory flow. *Clin Exp Allergy* 2000; 30:833-838
- Armitage P, Berry G. *Statistical Methods in Medical Research*. 2<sup>nd</sup> ed. Oxford:Ed. Blackwell Scientific Publications 1989
- Carlson JWK, Hoey H, Taylor MRH. Growth and other factors affecting peak expiratory flow rate. *Arch Dis Child* 1989; 64:96-102
- American Thoracic Society (ATS) Statement. Lung function testing of reference values and interpretative strategies. *Am Rev Respir Dis* 1991; 144:1202-1218
- European Respiratory Society (ERS). Lung volumes and forced ventilatory flows. Report Working Party on Standardization of Lung Function Test. *Eur Respir J* 1993; 6 :5-40
- Aivazis V, Hatzimichail A, Stavridis J, Bourli E , Konstantinidis Th , Katsougiannopoulos V. Growth and other factors affecting peak expiratory flow in Greek children. *Minerva Pediatrica* 2005; 57: 83-89
- Cotes JE, Leathart GL. Lung function throughout life: Determinants and reference values. *Lung Function Assessment and application in medicine*. 5<sup>th</sup> ed. Blackwell publications, 1993:445-513
- Paoletti P, Paggiaro PL, Lebowitz MD. Environmental factors in PEF variability. *Eur Respir J* 1997; 10:64-66
- Wihl JA, Malm L. Rhinomanometry and nasal peak expiratory and inspiratory flow rate. *Ann Allergy* 1988; 61:50-55
- Nathan Ra, Eccles R, Howarth PH, Steinsvag SK, Togios A. Objective monitoring of nasal patency and nasal physiology in rhinitis. *J Allergy Clin Immunol* 2005; 115: 442-459
- Wilson AM, Sims EJ, Robb F, Cockburn W, Lipworth BJ. Peak inspiratory flow rate is more sensitive than acoustic rhinomanometry or rhinomanometry in detecting corticosteroid response with nasal histamine challenge. *Rhinology* 2003; 41:16-20
- Trichopoulos D. *Medical Statistics. Basic Principles and methods of biomedical statistics*. Athens: Parisianos 1975
- Barbarousis D, Gioulekas D, Aivazis V, Kakavelas H. An investigation of the Lung Function of children and Teenagers of Thessaloniki. *Hell Rev Respir Dis* 1987; 5:223-236
- Mucha SM, de Tineo M, Naclerio RM, Baroody FM. Comparison of montelukast and pseudoephedrine in the treatment of allergic rhinitis. *Arch Otolaryngol Head Neck Surg* 2006; 132:164-172
- Barnes ML, Bialosterski BT, Gray RD, Fardon TC, Limorth BJ. Decongestant effects of nasal xylometazoline and mometasone furoate in persistent allergic rhinitis. *Rhinology* 2005; 43:291-295
- Oldenbeuving NB, Kleinjan A, Mulder PG, et al. Evaluation of an intranasal house dust mite provocation model as a tool in clinical research. *Allergy* 2005; 60:751-759
- Bhatia S, Baroody FM, de Tineo M, Naclerio RM. Increased nasal airflow with budesonide compared with desloratadine during the allergy season. *Arch Otolaryngol Head Neck Surg* 2005; 131:223-228
- Lund VJ, Black JH, Szabo LZ, Schrewelius C, Akerlund A. Efficacy and tolerability of budesonide aqueous nasal spray in chronic rhinosinusitis patients. *Rhinology* 2004; 42:57-62