

**Table 3.** Possible risk factors for poor outcome.

|                           | <b>All patients</b> | <b>Survivors</b> | <b>Non-survivors</b> | <b>p</b>     |
|---------------------------|---------------------|------------------|----------------------|--------------|
| Subjects                  | 105                 | 47               | 58                   | -            |
| MV Duration               |                     |                  |                      |              |
| <24h                      | 32/105              | 23/32            | 9/32                 | <b>0.005</b> |
| >24h                      | 73/105              | 24/73            | 49/73                |              |
| Vasopressors              |                     |                  |                      |              |
| Yes                       | 87/105              | 30/105           | 57/87                | <b>0.005</b> |
| No                        | 18/105              | 17/18            | 1/18                 |              |
| Associated organ failures |                     |                  |                      |              |
| <2                        |                     |                  |                      |              |
| >2                        | 53/105              | 45/53            | 8/53                 | <b>0.005</b> |
| Renal dysfunction         | 52/105              | 2/52             | 50/52                |              |
| Yes                       | 37/105              | 24/37            | 13/37                | 0.190        |
| No                        | 68/105              | 23/68            | 45/68                |              |
| Septic condition          |                     |                  |                      |              |
| Yes                       | 47/105              | 17/47            | 30/47                | <b>0.005</b> |
| No                        | 58/105              | 30/58            | 28/58                |              |