

Table 1: Univariate logistic regression analysis of risk factors for Crimean-Congo hemorrhagic fever virus (CCHFV) IgG seropositivity among the general human population in the area where the CCHFV strain AP92 had been isolated.

	Positive (%) N=6 (2.2)	Negative (%) N=271 (97.8)	p	OR	95% CI
Prefecture			0.504	1.73	0.34, 8.77
Pella	3 (2.9)	99 (97.1)			
Imathia	3 (1.7)	172 (98.3)		reference	
Median age (y)	67	47	0.043	1.06	1.00, 1.13
Sex			0.221	3.86	0.45, 33.45
Male	5 (3.2)	153 (96.8)			
Female	1 (0.8)	118 (99.2)		reference	
Occupation			0.579	1.84	0.21, 16.07
Farmer	5 (2.5)	195 (97.5)			
Other	1 (1.4)	72 (98.6)		reference	
Tick bite			0.040	13.00	1.31, 149.46
Yes	2 (8.3)	22 (91.7)			
No	1 (0.7)	143 (99.3)		reference	
Missing data	3	106			
Altitude			0.025		
0-80 m	0 (0)	77 (100)			
81-300 m	5 (5.9)	80 (94.1)			
>300 m	1 (1.0)	104 (99.0)			

OR: odds ratio. CI: confidence intervals. y: years. m: meters.