

Table 1b: Concomitant interacting medications and comorbidity of the 101 warfarin-treated patients included in the study.

Concomitant interacting medications^a	
Antiepileptics (phenytoin, levetiracetam, gabapentin, ect.)	5 (4.95%)
Antibiotics (quinolones)	2 (1.98%)
Lipid-lowering agents (statins)	36 (35.64%)
H ₂ blockers (famotidine)	1 (0.99%)
Proton pump inhibitors	7 (6.93%)
Angiotensin II receptor antagonists	14 (13.86%)
β-blockers	69 (68.32%)
Calcium Channel Blockers	11 (10.89%)
α-blockers	11 (10.89%)
Diuretics	57 (56.44%)
Bronchodilators	8 (7.92%)
Antiagregants (ASA, etc.)	57 (56.44%)
Oral iron therapy	3 (2.97 %)
Antiischemic drugs	7 (6.93%)
Coronary vasodilators	9 (8.91%)
Antidiabetic drugs	13 (12.87%)
Thyroid drugs	7 (6.93%)
Antidepressant drugs	6 (5.94%)
Other	11 (10.89%)
Comorbidity	
Hypertension	60 (59.40 %)
Arrhythmia	1 (0.99 %)
Type II diabetes	18 (17.82 %)

Data are presented as number of patients (%), ^a: patients might have used more than one concomitant interacting medication.