

Table 2. Distribution of animals treated with Tamsulosin (Group 2) by weight, intervention time, operated eye, billowing grade during cannula irrigation and irrigation aspiration.

| Group 2 - tamsulosin 0.4 mg/kg | | | | | | | |
|--------------------------------|--|------------------------------|---------------|---------------------------------------|--|----------------|----------------------------------|
| Animal No. | Weight (kg) Mean \pm SD 4.48 \pm 0.913 | Intervention (date, time) | Eye | Billowing grade cannula irrigation | Billowing grade irrigation aspiration | Total grade | Remarks |
| T1 | 4.5 | 29.05.12 14:50 | Left Right | 0 0 | 0 0 | 0 0 | |
| T2 | 4.4 | 29.05.12 17:40 | Left Right | 3 2 | 2 1 | 5 3 | |
| T3 | 4.8 | 30.05.12 14:02 | Left Right | 2 3 | 1 1 | 3 4 | |
| T4 | 4.8 | 29.05.12 16:11 | Left Right | 2 2 | 1 1 | 3 3 | |
| T5 | 4.2 | 29.05.12 15:55 | Left Right | - - | - - | - - | incomplete protocol |
| T6 | 4 | 29.05.12 18:10 | Left Right | 1 0 | 0 0 | 1 0 | |
| T7 | 4 | 29.05.12 15:45 | Left Right | - - | - - | - - | bilateral incomplete protocol |
| T8 | 3.5 | 30.05.12 14:02 | Left Right | 1 1 | 0 0 | 1 1 | |
| T9 | 3.6 | 30.05.12 14:02 | Left Right | - 3 | - 1 | - 4 | Left eye - incorect wound |
| T10 | 3.6 | 30.05.12 14:02 | Left Right | - 2 | - 2 | - 4 | Left eye - incorect wound |

SD: standard deviation.