

**Table 3.** Distribution of animals treated with finasteride (Group 3) by weight, intervention time, operated eye, billowing grade during cannula irrigation and irrigation aspiration.**Group 3 – Finasteride 2.5 mg/kg**

Animal No.	Weight (kg) Mean $\pm$ SD 4.48 $\pm$ 0.913	Intervention (date, time)	Eye	Billowing grade cannula irrigation	Billowing grade irrigation aspiration	Total grade	Remarks
F1	3.5	29.05.12 17:55	Left Right	1 2	0 1	1 3	
F2	3.8	29.05.12 17:00	Left Right	1 2	0 1	1 3	
F3	4.4	29.05.12 16:25	Left Right	0 2	0 1	0 3	
F4	5	29.05.12 16:45	Left Right	2 1	0 0	2 1	
F5	4.2	29.05.12 18:10	Left Right	3 3	2 2	5 5	
F6	4.8	30.05.12 14:10	Left Right	3 3	2 2	5 5	
F7	4.4	29.05.12 18:30	Left Right	2 1	1 1	3 2	
F8	3.5	29.05.12 17:48	Left Right	1 1	1 1	2 2	
F9	4.4	29.05.12 18:19	Left Right	2 3	1 1	3 4	
F10	4.4	30.05.12 15:00	Left Right	0 0	0 0	0 0	Increased dosage of anesthetic drug

SD: standard deviation.