

Table 2: Laboratory findings of the study population consisting of 199 patients, of whom 103 non-dipper and 96 dipper.

| Variables | All patients (n =199) | Non-dipper (n =103) | Dipper (n =96) | p value |
|-----------------------------------|-----------------------|---------------------|--------------------|--------------|
| Fasting glucose (mg/dl) | 89.6 ± 6.4 | 90.1 ± 5.5 | 89.2 ± 7.1 | 0.414 |
| Creatinine (mg/dl) | 0.8 ± 0.1 | 0.8 ± 0.1 | 0.8 ± 0.2 | 0.350 |
| Albumin (mg/dl) | 46.4 ± 2.6 | 46.2 ± 2.4 | 46.6 ± 2.7 | 0.314 |
| Triglyceride (mg/dl) | 141 (57-735) | 144 (57-735) | 125 (64-360) | 0.123 |
| LDL cholesterol (mg/dl) | 125.9 ± 16.3 | 124.1 ± 16.8 | 127.8 ± 15.5 | 0.115 |
| HDL cholesterol (mg/dl) | 49.7 ± 13.2 | 49.1 ± 13.5 | 50.5 ± 12.9 | 0.470 |
| ESR (mm/hour) | 14.4 ± 3.5 | 14.5 ± 3.7 | 13.8 ± 3.3 | 0.177 |
| CRP (mg/dl) | 3.1 (0.3-12) | 3.5 (0.4-12) | 2.7 (0.3-8) | 0.068 |
| Proteinuria (mg/day) | 93 (6-1949) | 95.4 (13.9-1949) | 90.8 (6-1619.5) | 0.466 |
| WBC (x10 ³ /μL) | 7.0 ± 1.4 | 6.9 ± 1.4 | 7.2 ± 1.3 | 0.046 |
| Lymphocyte (x10 ³ /μL) | 2.2 (1-4) | 2,1 (1-3,9) | 2,2 (1-4) | 0,001 |
| Platelet (x10 ³ /μL) | 279.7 ± 57.4 | 282 ± 60.6 | 277.2 ± 54 | 0.559 |
| PLR | 124 (53.5-400.0) | 132.2 (69.6-400) | 117.0 (53.5-192.3) | 0.001 |
| Hemoglobin (g/dl) | 13.9 ± 1.4 | 13.8 ± 1.4 | 14.1 ± 1.4 | 0.109 |

Data are presented as the number of patients or mean value ± standard deviation, p <0.05 is considered statistically significant for all tests, n: number, LDL: low-density lipoprotein, HDL: high-density lipoprotein, ESR: erythrocyte sedimentation rate, CRP: C-reactive protein, WBC: White blood cell, PLR: Platelet-to-lymphocyte ratio, SD: standard deviation.