

Table 1: Demographics, hemodialysis parameters, comorbidity and therapeutic regimen of the 48 hemodialysis patients who participated in the study.

Demographics	Patients (n=48)
Sex (Male/Female)	25/23
Age (years)	56.9 ± 9.3
Therapy Duration (months)	43 ± 31
Residual Diuresis (ml/day)	n=15, 160 ± 50
Renal Replacement Therapy Parameters	Patients (n=48)
Mode (Hemodialysis/Hemodiafiltration)	31/17
Vascular Access (Fistula/Graft/Catheter)	32/4/12
Dialysis Membrane (Polysulfone/Polyethersulfone)	26/22
Membrane Surface (m ²)	1.7 ± 0.1
Dialysis Anticoagulation-Low Molecular Weight Heparin (IU)	4,300 ± 1,800
Comorbidity	Patients (n=48)
Diabetes Melitus	40
Coronary Heart Disease	8
Arterial Hypertension	32
Peripheral Vascular Disease	11
Pulmonary Disease	2
Renal Transplantation	4
Drug Therapy	Patients (n=48)
B-blockers	19
Statins/Ω ₃ Fatty Acids	25/13
Angiotensin Converting Enzyme inhibitors/ Angiotensin II Receptor blockers	19
Vitamin D (Alphacalcidol/Paracalcitol) (µg/week)	14 (1µg) / 5 (7.5µg)
Cinacalcet (mg/day)	9 (15 ± 22.5mg)
Epoetin β (IU/week)	14,440 ± 8,370
Iron Sucrose (mg/week)	143 ± 83
Vitamin B complex (B ₁ , B ₆ , B ₁₂)	48
Carnitine	48
Folate	31

All values are mean ± standard deviation.