

**Table 4:** Demographic features in the 47 patients with cardiothoracic index >0.5 and in the 53 with cardiothoracic index ≤0.5.

Variables	CTI >0.50 (n=47)	CTI ≤0.50 (n=53)	p
Age (years)	63.40 ± 11.9	56.95 ± 13.2	0.010
Sex (male) n (%)	34 (61.8)	21 (46.7)	0.130
Height (m)	1.62 ± 0.09	1.60 ± 0.09	0.341
Weight (kg)	67.37 ± 14.8	66.20 ± 14.7	0.868
BMI(kg/m <sup>2</sup> )	26.03 ± 5.5	25.37 ± 4.8	0.444
Body surface area (m <sup>2</sup> )	1.73 ± 0.2	1.72 ± 0.2	0.905
HD duration (year)	7 (1-31)	8 (1-23)	0.622
Diabetes mellitus n (%)	12 (21.8)	16 (35.6)	0.179
Antihypertensive treatment	23 (41.8)	18 (40.0)	0.995
Vascular access n (%)			
Fistula	49 (89.1)	37 (82.2)	0.24
Catheter	5 (9.1)	8 (17.8)	
Graft	1 (1.8)		

CTI: cardiothoracic index, BMI: body mass index, HD: hemodialysis, n: number, data are presented as mean ± standard deviation or number (percentage).