

Table 5: The correlation between the duration of T2DM and absolute and inter-wave latencies of BAEP.

BAEP latencies	Duration of diabetes		
	≤ 10 years (n = 51) Mean ± SD (ms)	>10 years (n = 29) Mean ± SD (ms)	p value
I R	1.34 ± 0.04	1.34 ± 0.05	0.995
I L	1.33 ± 0.04	1.35 ± 0.04	0.261
III R	3.60 ± 0.08	3.62 ± 0.08	0.300
III L	3.61 ± 0.08	3.61 ± 0.09	0.686
V R	5.47 ± 0.10	5.51 ± 0.08	0.091
V L	5.47 ± 0.10	5.51 ± 0.08	0.119
I-III R	2.26 ± 0.07	2.27 ± 0.08	0.462
I-III L	2.26 ± 0.07	2.26 ± 0.07	0.844
III-V R	1.86 ± 0.07	1.89 ± 0.04	0.127
III-V L	1.86 ± 0.06	1.88 ± 0.05	0.087
I-V R	4.13 ± 0.98	4.17 ± 0.07	0.066
I-V L	4.13 ± 0.08	4.16 ± 0.06	0.167

T2DM: Type 2 diabetes mellitus, n: number of subjects, BAEP: brainstem auditory evoked potentials, SD: standard deviation, ms: millisecond, R: right ear, L: left ear, p: Student's t-test value (p <0.05, bolded if significant).