

### References of the supplementary Tables

- Bischof DB, Ganter MT, Shore-Lesserson L, Hartnack S, Klaghofer R, Graves K, et al. Viscoelastic blood coagulation measurement with Sonoclot predicts postoperative bleeding in cardiac surgery after heparin reversal. *J Cardiothorac Vasc Anesth.* 2015; 29: 715-722.
- Greilich PE, Edson E, Rutland L, Jessen ME, Key NS, Levy JH, et al. Protocol adherence when managing massive bleeding following complex cardiac surgery: a study design pilot. *J Cardiothorac Vasc Anesth.* 2015; 29: 303-310.
- Doussau A, Perez P, Puntous M, Calderon J, Jeanne M, Germain C, et al; PLASMACARD Study Group. Fresh-frozen plasma transfusion did not reduce 30-day mortality in patients undergoing cardiopulmonary bypass cardiac surgery with excessive bleeding: the PLASMACARD multicenter cohort study. *Transfusion.* 2014; 54: 1114-1124.
- Tanaka KA, Egan K, Szlam F, Ogawa S, Roback JD, Sreeram G, et al. Transfusion and hematologic variables after fibrinogen or platelet transfusion in valve replacement surgery: preliminary data of purified lyophilized human fibrinogen concentrate versus conventional transfusion. *Transfusion.* 2014; 54: 109-118.
- Kremke M, Tang M, Bak M, Kristensen KL, Hindsholm K, Andreassen JJ, et al. Antiplatelet therapy at the time of coronary artery bypass grafting: a multicentre cohort study. *Eur J Cardiothorac Surg.* 2013; 44: e133-e140.
- Andersen ND, Bhattacharya SD, Williams JB, Fosbol EL, Lockhart EL, Patel MB, et al. Intraoperative use of low-dose recombinant activated factor VII during thoracic aortic operations. *Ann Thorac Surg.* 2012; 93: 1921-1928; discussion 1928-1929.
- Chapman AJ, Blount AL, Davis AT, Hooker RL. Recombinant factor VIIa (NovoSeven RT) use in high risk cardiac surgery. *Eur J Cardiothorac Surg.* 2011; 40: 1314-1318; discussion 1318-1319.
- Williams JB, Phillips-Bute B, Bhattacharya SD, Shah AA, Andersen ND, Altintas B, et al. Predictors of massive transfusion with thoracic aortic procedures involving deep hypothermic circulatory arrest. *J Thorac Cardiovasc Surg.* 2011; 141: 1283-1288.
- Girdauskas E, Kempfert J, Kuntze T, Borger MA, Enders J, Fassl J, et al. Thromboelastometrically guided transfusion protocol during aortic surgery with circulatory arrest: a prospective, randomized trial. *J Thorac Cardiovasc Surg.* 2010; 140: 1117-1124.e2.
- Willis C, Bird R, Mullany D, Cameron P, Phillips L. Use of rFVIIa for critical bleeding in cardiac surgery: dose variation and patient outcomes. *Vox Sang.* 2010; 98: 531-537.
- Christensen MC, Krapf S, Kempel A, von Heymann C. Costs of excessive postoperative hemorrhage in cardiac surgery. *J Thorac Cardiovasc Surg.* 2009; 138: 687-693.
- Masud F, Bostan F, Chi E, Pass SE, Samir H, Stuebing K, et al. Recombinant factor VIIa treatment of severe bleeding in cardiac surgery patients: a retrospective analysis of dosing, efficacy, and safety outcomes. *J Cardiothorac Vasc Anesth.* 2009; 23: 28-33.
- Wasowicz M, Meineri M, McCluskey SM, Mitsakakis N, Karkouti K. The utility of thromboelastography for guiding recombinant activated factor VII therapy for refractory hemorrhage after cardiac surgery. *J Cardiothorac Vasc Anesth.* 2009; 23: 828-834.
- Karkouti K, Beattie WS, Arellano R, Aye T, Bussières JS, Callum JL, et al. Comprehensive Canadian review of the off-label use of recombinant activated factor VII in cardiac surgery. *Circulation.* 2008; 118: 331-338.
- Trowbridge C, Stammers A, Klayman M, Brindisi N, Woods E. Characteristics of uncontrolled hemorrhage in cardiac surgery. *J Extra Corpor Technol.* 2008; 40: 89-93.
- Karkouti K, O'Farrell R, Yau TM, Beattie WS; Reducing Bleeding in Cardiac Surgery Research Group. Prediction of massive blood transfusion in cardiac surgery. *Can J Anaesth.* 2006; 53: 781-794.
- Chen L, Bracey AW, Radovancevic R, Cooper JR Jr, Collard CD, Vaughn WK, et al. Clopidogrel and bleeding in patients undergoing elective coronary artery bypass grafting. *J Thorac Cardiovasc Surg.* 2004; 128: 425-431.
- Stein P, Bosshart M, Brand B, Schlicker A, Spahn DR, Bettex D. Dabigatran anticoagulation and Stanford type A aortic dissection: lethal coincidence: Case report with literature review. *Acta Anaesthesiol Scand.* 2014; 58: 630-637.
- Warkentin TE, Margetts P, Connolly SJ, Lamy A, Ricci C, Eikelboom JW. Recombinant factor VIIa (rFVIIa) and hemodialysis to manage massive dabigatran-associated postcardiac surgery bleeding. *Blood.* 2012; 119: 2172-2174.
- Barua A, Rao VP, Ramesh B, Barua B, El-Shafei H. Salvage use of activated recombinant factor VII in the management of refractory bleeding following cardiac surgery. *J Blood Med.* 2011; 2: 131-134.
- Bishop CV, Renwick WE, Hogan C, Haeusler M, Tuckfield A, Tatoulis J. Recombinant activated factor VII: treating postoperative hemorrhage in cardiac surgery. *Ann Thorac Surg.* 2006; 81: 875-879.
- van de Garde EM, Bras LJ, Heijmen RH, Knibbe CA, van Dongen EP, Wiltink EH, et al. Low-dose recombinant factor VIIa in the management of uncontrolled postoperative hemorrhage in cardiac surgery patients. *J Cardiothorac Vasc Anesth.* 2006; 20: 573-575.
- Goudie R, Sterne JAC, Verheyden V, Bhabra M, Ranucci M, Murphy GJ. Risk scores to facilitate preoperative prediction of transfusion and large volume blood transfusion associated with adult cardiac surgery. *Br J Anaesth.* 2015; 114: 757-766.
- Vuylsteke A, Pagel C, Gerrard C, Reddy B, Nashef S, Aldam P, et al. The Papworth Bleeding Risk Score: a stratification scheme for identifying cardiac surgery patients at risk of excessive early postoperative bleeding. *Eur J Cardiothorac Surg.* 2011; 39: 924-930.
- Ranucci M, Castelvechchio S, Frigiola A, Scolletta S, Giomarelli P, Biagioli B. Predicting transfusions in cardiac surgery: the easier, the better: the Transfusion Risk and Clinical Knowledge score. *Vox Sang.* 2009; 96: 324-332.
- Alghamdi AA, Davis A, Brister S, Corey P, Logan A. Development and validation of Transfusion Risk Understanding Scoring Tool (TRUST) to stratify cardiac surgery patients according to their blood transfusion needs. *Transfusion.* 2006; 46: 1120-1129.
- Litmathe J, Boeken U, Feindt P, Gams E. Predictors of homologous blood transfusion for patients undergoing open heart surgery. *Thorac Cardiovasc Surg.* 2003; 51: 17-21.
- Magovern JA, Sakert T, Benckart DH, Burkholder JA, Liebler GA, Magovern GJ Sr, et al. A model for predicting transfusion after coronary artery bypass grafting. *Ann Thorac Surg.* 1996; 61: 27-32.
- Dyke C, Aronson S, Dietrich W, Hofmann A, Karkouti K, Levi M, et al. Universal definition of perioperative bleeding in adult cardiac surgery. *J Thorac Cardiovasc Surg.* 2014; 147: 1458-1463.e1.
- Karkouti K, Wijesundera DN, Yau TM, Beattie WS, Abdelnaem E, McCluskey SA, et al. The independent association of massive blood loss with mortality in cardiac surgery. *Transfusion.* 2004; 44: 1453-1462.