

**Table 2:** Clinical characteristics of the 49 patients with Crohn's disease (CD) who were divided into two groups: 24 patients with active CD (ACDP) and 25 patients with inactive CD (ICDP).

	ACDP (n =24)	ICDP (n =25)
Location of disease		
Ileitis	10 (45.5 %)	15 (60 %)
Colitis	4 (18.1 %)	2 (8 %)
Ileocolitis	8 (36.4 %)	8 (32 %)
Extraintestinal manifestations		
No	22 (91.7 %)	21 (84 %)
Arthritis	2 (8.3 %)	1 (4 %)
Ankylosing spondylitis	-	1 (4 %)
Sacroileitis	-	2 (8 %)
Behavior of disease		
Non-stricturing non-penetrating	9 (37.5 %)	16 (64 %)
Stricturing	9 (37.5 %)	5 (20 %)
Penetrating	6 (25 %)	4 (16 %)
Medications*		
Azathioprine	14 (58.3 %)	13 (52 %)
5-Aminosalicylic acid	10 (41.7 %)	4 (16 %)
Corticosteroids	13 (54.2 %)	10 (40 %)
TNF- $\alpha$ blocker	2 (8.3 %)	7 (28 %)
No medications	5 (20.8 %)	-

Values are expressed as numbers (% or range in brackets), ACDP: patients with active Crohn's disease, ICDP: patients with inactive Crohn's disease, \*: Some patients were taking more than one medication at the same time.