

Table 1: Demographics, comorbidities and source of infection in the whole sample, and comparison of the variables in Group I (patients with no contact with a healthcare facility for more than 48 hours within the preceding six months) and Group II (patients admitted from another healthcare facility/nursing home or mentioned contact with a healthcare facility for more than 48 hours within the preceding six months).

	Total	Group I	Group II	p
Number of patients (%)	288 (100.0)	181 (62.8)	107 (37.2)	
Age (years)	82 [60-97]	80.4	80.3	0.885
Sex (male) (%)	42.7	43.1	42.0	0.863
Diabetes Mellitus (%)	31.9	30.9	33.7	0.634
Heart Failure (%)	53.4	51.7	56.1	0.478
Respiratory Failure (%)	19.5	19.9	18.9	0.833
Neurologic Deficits (%)	33.8	30.4	39.6	0.110
Chronic Kidney Disease (%)	57.7	57.5	55.8	0.625
Acute Kidney Injury (%)	36.9	35.7	38.8	0.599
Pneumonia (%)	54.4	58.7	49.1	0.058
Urinary Tract Infection (%)	42.5	33.9	56.8	<0.001
Other (infection) (%)	27.0	23.4	33.0	0.082
Bacteremia (%)	15.3	11.1	22.4	0.009
Outcome (Death) (%)	21.5	18.2	27.1	0.077
Duration of stay (survivors) (days)	8.6 [1-38]	8	9.7	0.050