

Table 5: Logistic regression (FACTOR A derived from factor analysis and available algorithms for hepatic steatosis prediction).

		B	S.E.	Wald	df	Sig.	Exp (B)
<i>Model 1</i>	FACTOR A	1.455	0.433	11.292	1	0.001	4.283
	Constant	0.376	0.271	1.918	1	0.166	1.456
<i>Model 2</i>	FLI	0.031	0.009	11.491	1	0.001	1.032
	Constant	-1.764	0.663	7.081	1	0.008	0.171
<i>Model 3</i>	HSI	0.107	0.043	6.137	1	0.013	1.112
	Constant	-3.984	1.713	5.411	1	0.020	0.019
<i>Model 4</i>	TyG	0.652	0.436	2.240	1	0.135	1.920
	Constant	-5.459	3.840	2.022	1	0.155	0.004
<i>Model 5</i>	LAP	0.005	0.004	1.124	1	0.289	1.005
	Constant	-0.080	0.409	.039	1	0.844	0.923
<i>Model 6</i>	VAI	0.022	0.089	0.063	1	0.802	1.023
	Constant	0.225	0.338	0.444	1	0.505	1.253

FLI: fatty liver index, HSI: hepatic steatosis index, LAP: lipid accumulation product, VAI: visceral adiposity index, TyG: triglyceride and glucose index, B: coefficient, S.E.: standard error, Wald: Wald, df: degree of freedom, Sig: significance, Exp (B): exponential B coefficient.