

Table 1: Demographic, clinical and analytical data of the 87 patients enrolled in this cross-sectional and observational study.

| | |
|---|-------------------|
| Demographics | |
| Age (years) | 52.3 ± 12.7 |
| Sex (M/F) | 40/47 |
| BMI (kg/m ²) | 30.16 ± 7.20 |
| Obese patients (BMI >30 kg/m ²) | 26 of 87 patients |
| Diabetes (yes/no) | 80/7 |
| Hypertension (yes/no) | 42/45 |
| Cardiovascular disease (yes/no) | 17/70 |
| rhEPO therapy (yes/no) | 42/45 |
| Modality of peritoneal dialysis | |
| CAPD/APD (%) | 90.8/9.2 |
| Icodextrin | 44 of 87 patients |
| Aminoacid-based solutions | 18 of 87 patients |
| Analytical data | |
| Albumin (g/dl) | 3.43 ± 0.51 |
| Hemoglobin (g/dl) | 10.6 ± 1.6 |
| Dialysis parameters | |
| Duration(months) | 48 |
| Weekly Kt/V | 2.63±1.16 |

Values are given in mean ± standard deviation, unless otherwise stated in brackets, CAPD: continuous ambulatory peritoneal dialysis, APD: automated peritoneal dialysis, Kt/V: urea clearance.