

Table 3: Comparisons of body mass index, Beck Depression Inventory, residual renal function and parameters of peritoneal equilibration test in patients using and not using icodextrin-based solutions.

Parameters	Patients using icodextrin -based solutions (n: 44)	Patients not using icodextrin -based solutions (n:43)	p value
BMI	29.65 (25.42-33.90)	28.80 (25-31)	0.33
BECK	10 (6-17)	9 (3-18)	0.81
Kt/V	2.23 (2-3.13)	2.47 (1.82-2.92)	0.39
nPCR (gr/kg/day)	1.06 (0.81- 1.68)	2.04 (1.26-2.5)	<0.001
Creatinine clearance (ml/min)	62.65 (50-98)	71.83 (49-110)	0.39
Residual GFR (ml/min)	6.75 (1.2-24)	10.97 (0.3-35)	0.76
Residual urine volume (cc)	550 (25- 1181)	700 (300-1400)	0.20
D/P crt	0.64 ± 0.11	0.65 ± 0.12	0.57

n: Number, Values are expressed as median (interquartile range) or mean ± standard deviation, BMI: body mass index, BECK: Beck Depression Inventory, Kt/V: urea clearance, nPCR: normalized protein catabolic rate, D/P crt: dialysate-peritoneal creatinine ratio.