

Table 2: Distribution of *cytochrome P450 2E1 (CYP2E1)* RsaI/PstI genotype among alcoholic pancreatitis cases and controls included in the meta-analysis.

First author	Year	Type of AP	Controls				Cases			Controls		Cases	
			c1/c1	c1/c2	c2/c2	HWE (p value)	c1/c1	c1/c2	c2/c2	c1	c2	c1	c2
Matsumoto ²⁸	1996	CAP	39	21	2	>0.05	9	2	0	99	25	20	2
Chao ²⁵	1997	AAP	68	43	8	>0.05	30	15	3	179	59	75	21
Maruyama ³³	1999	CAP	30	15	1	>0.05	30	21	2	75	17	81	25
Yang ³¹	2001	CAP	192	9	0	>0.05	37	1	0	393	9	75	1
Frenzer ²⁶	2002	AAP	192	9	0	>0.05	18	1	0	393	9	37	1
		CAP	242	15	0	>0.05	65	6	0	499	15	136	6
Burim ³⁰	2004	CAP	234	27	1	>0.05	11	3	0	495	29	25	3
Kim ²⁷	2004	NM	51	34	15	>0.05	17	11	1	136	64	45	13
Verlaan ²⁹	2004	CAP	122	6	0	>0.05	75	7	0	250	6	157	7
Cichoz-Lach ²²	2008	CAP	97	0	0	>0.05	42	2	0	194	0	86	2
Gubergrits ³²	2014	CAP	55	22	3	>0.05	36	26	10	132	28	98	46

HWE: Hardy-Weinberg Equilibrium, AP: alcoholic pancreatitis, CAP: chronic alcoholic pancreatitis, AAP: acute alcoholic pancreatitis, NM: not mentioned.