

**Table 3:** Dermatological findings in the study (hemodialysis, peritoneal dialysis, kidney transplant, chronic kidney disease, and healthy control) groups.

Dermatological findings	Group 1	Group 2	Group 3	Group 4	Group 5	p-value
Pruritus	10 (33.3)	10 (38.5) <sup>a</sup>	2 (6.7) <sup>b</sup>	8 (26.7)	3 (10)	<b>0.01</b>
Xerosis	25 (83.3) <sup>c</sup>	12 (46.2)	9 (30)	13 (44.8)	1 (3.3) <sup>d</sup>	<b>0.0001</b>
Hyperpigmentation	13 (43.3)	13 (50)	13 (43.3)	11 (37.9)	1 (3.3) <sup>e</sup>	<b>0.001</b>
Alopecia	16 (53.3) <sup>f</sup>	11 (42.3) <sup>g</sup>	7 (23.3)	5 (17.2)	4 (13.3)	<b>0.003</b>

Values are presented as the number of patients and percentage in brackets. Group 1: hemodialysis, group 2: peritoneal dialysis, group 3: kidney transplant, group 4: chronic kidney disease, group 5: healthy control. <sup>a</sup>: group 2 versus group 5, <sup>b</sup>: group 3 versus groups 1,2, and 4, <sup>c</sup>: group 1 versus groups 2,3, and 4, <sup>d</sup>: group 5 versus groups 1,2,3, and 4, <sup>e</sup>: group 5 versus groups 1,2,3, and 4, <sup>f</sup>: group 1 versus groups 3,4, and 5, <sup>g</sup>: group 2 versus group 5.