

Table 8: Comparisons of serum interleukin 31 (IL-31) levels (pg/mL) according to dermatological findings in the study (hemodialysis, peritoneal dialysis, kidney transplant, chronic kidney disease, and healthy control) groups.

Finding		Group 1	Group 2	Group 3	Group 4	Group 5
Longitudinal ridges	present	0.43 (0.01-5.08)	0.28 (0.01-0.89)	0.02 (0.01-3.17)	0.92 (0.46-2.29)	1.02 (0.02-11.53)
	absent	0.38 (0.01-3.92)	0.48 (0.01-3.25)	1.25 (0.01-2.63)	0.5 (0.04-2.52)	0.77 (0.01-2.44)
	p-value	0.8	0.31	0.63	0.02	0.3
Tinea pedis	present	0.81 (0.01-1.62)	0.35 (0.02-0.49)	0.006 (0.01-1.19)	0.64 (0.09-2.52)	1.45 (0.06-11.53)
	absent	0.43 (0.01-5.08)	0.48 (0.01-3.25)	1.42 (0.01-3.17)	0.55 (0.04-2.1)	0.87 (0.01-2.04)
	p-value	0.83	0.42	0.01	0.4	0.21

Values of IL-31 are presented median and minimum-maximum in brackets. Group 1: hemodialysis, group 2: peritoneal dialysis, group 3: kidney transplant, group 4: chronic kidney disease, group 5: healthy control. CKD: chronic kidney disease. IL-31: interleukin 31.