

**Table 9:** Comparisons of serum uridine diphosphate glucose ceramide glucosyltransferase (ng/mL) according to dermatological findings in the study (hemodialysis, peritoneal dialysis, kidney transplant, chronic kidney disease, and healthy control) groups.

Finding		Group 1	Group 2	Group 3	Group 4	Group 5
Pruritus	present	18.3 (9.3-45.4)	15.7 (1-29.4)	19 (13-25.2)	13.5 (8.7-22)	39.8 (16-41)
	absent	20.3 (6.4-37.7)	25.4 (5.1-42.7)	16.4 (5.9-39.5)	15.4 (8.8-40.5)	15.5 (7-41)
	p-value	0.79	<b>0.045</b>	0.86	0.18	0.06
Xerosis	present	19.6(6.3-45.4)	17.1(5.1-40.9)	25.2(8.1-39)	13.8(8.7-17.1)	20.4(20.4)
	absent	16.1(8.4-30.8)	24.5(1-42.7)	16.3(5.8-39.5)	16.7(10-40.5)	16(7-41)
	p-value	0.35	0.3	0.23	<b>0.01</b>	0.45
Splinter hemorrhages	present	18.3 (14.2-24.3)	22.9 (6.5-40.9)	12.5 (5.8-17.9)	14.6 (14.5-14.6)	
	absent	19.6(6.3-45.4)	22.2(1-42.7)	19.1(7.8-39.5)	15.1(8.7-40.5)	16.3(7-41)
	p-value	0.71	0.88	<b>0.02</b>	0.93	

Values of UGCG are presented median and minimum-maximum in brackets. Group 1: hemodialysis, group 2: peritoneal dialysis, group 3: kidney transplant, group 4: chronic kidney disease, group 5: healthy control. UGCG: uridine diphosphate glucose ceramide glucosyltransferase.