

Table 5: Delay rates for at least one dose of recommended vaccines for public and private nurseries, and for Greek and non-Greek families. Data are expressed as percentages with correspondent numbers of children in parentheses [% (n)].

Vaccine	Public nurseries (n =187)	Private nurseries (n =245)	p (public vs private)	Greek families (n =380)	Non-Greek families (n =52)	p (Greek vs non-Greek)
DTaP-IPV	52.4 (98)	46.5 (114)	0.24	45.7 (174)	65.3 (34)	0.03
Hib	54.5 (102)	48.6 (119)	0.21	49.2 (187)	65.4 (35)	0.03
HepB	48.1 (90)	38.5 (88)	0.01	41.6 (158)	52.0 (27)	0.16
PCV	80.7 (151)	72.6 (178)	0.05	76.8 (292)	71.1 (37)	0.45
MCC (2011)	73.8 (138)	54.2 (133)	<0.001	63.2 (240)	53.8 (28)	0.6
MCC (2015)	35.8 (67)	41.2 (101)	0.31	38.4 (146)	44.2 (23)	0.56
MMR	27.3 (51)	20.8 (51)	0.12	24.7 (94)	17.3 (9))	0.18
VAR	47.1 (88)	47.7 (117)	0.89	46.6 (177)	53.8 (28)	0.34
HepA	92.5 (173)	89.4 (219)	0.34	89.7 (341)	98 (51)	0.15
Rota	59.8 (112)	53.8 (132)	0.47	56.8 (216)	53.8 (28)	0.89

Reported p values are for comparisons between public versus private nurseries and Greek versus non-Greek families. DTaP-IPV: diphtheria-tetanus-acellular pertussis-inactivated polio, Hib: *Haemophilus influenzae* type b, HepB: hepatitis B, PCV: conjugated pneumococcal, MCC: conjugated meningococcal serogroup C, MMR: measles-mumps-rubella, VAR: varicella, HepA: hepatitis A, Rota: rotavirus.