Surgically Therapy Group (OG) Conservative Therapy Group (NOG)

Mean ± SD 95 % CI Mean ± SD 95 % CI

p value

1.25-1.85

5.58-35.17

17.82-91.25

6.49-9.58

OG: operated case group, NOG: Non-operatively treated case group, SD: standard deviation, CI: confidence interval, WBC: white blood cell,

 $9.06 \pm 4.83$ 

 $1.6 \pm 0.87$ 

 $12.45 \pm 5.68$ 

 $59.59 \pm 70.91$ 

 $7.21 \pm 5.60$ 

7.45-10.68

1.31-1.90

3.54-32.12

35.94-83.23

5.38-9.08

0.101\*

0.701\*\*

0.225\*\*

0.670\*\*

0.023\*\*

 Mean  $\pm$  SD
 95 % CI

 Neutrophil ( $\times 10^9$  cells/L)
 12.0  $\pm$  4.60
 4.58-10.68

Lymphocyte (×10<sup>9</sup> cells/L)

**WBC** ( $\times 10^9$  cells/L)

CRP (mg/L)

**NLR** 

**Table 1:** Comparison of laboratory parameters of surgical and conservative therapy groups.

 $1.5 \pm 0.56$ 

 $14.04 \pm 6.58$ 

 $54.54 \pm 68.89$ 

 $8.03 \pm 2.89$ 

CRP: C-reactive protein, NLR: neutrophil-to-lymphocyte ratio, \*: t-test, \*\*: Mann-Whitney U test.