

Table 2: Baseline characteristics and failed treatments of the reported eight patients with hidradenitis suppurativa complicated with amyloidosis.

Patient	Age/Gender	Smoking Status	Comorbidities	Duration of HS	Hurley Stage	Previous Treatments
1	50/M	Former	FMF + arthritis	20	3	Colchicine + Ab + Surgery + NSAII + GC + Dapson + Apremilast
2	39/F	Former	-	11	3	Ab + acitresin + isotretinoin + GC + surgery + ADA
3	44/M	Active	-	18	2-3	Ab + acitresin + surgery + ADA
4	62/M	Active	Hepatitis B COPD	30	1-2	Ab + GC
5	43/M	Active	-	15	2-3	Ab + GC + Surgery
6	65/M	Former	DM, CAD	10	3	Ab + isotretinoin + GC + surgery
7	44/F	Active	-	16	3	Ab + acitresin + isotretinoin + GC + surgery + IFX
8	35/M	Never	Crohn	7	2-3	Ab + GC + MTX + Aza + ET

M: male, F: female, Ab: systemic antibiotics, FMF: Familial Mediterranean Fever, COPD: chronic obstructive pulmonary disease, DM: diabetes, CAD: coronary artery disease, ADA: adalimumab, GC: glucocorticoid, iGC: intralesional glucocorticoids, IFX: Infliximab, MTX: methotrexate, Aza: azathioprine, ET: etanercept.