Table 3: Follow-up of the reported eight patients with hHidradenitis suppurativa complicated with amyloidosis.

atient	Laboratory results at	Biopsy site	Follow up (months)	Intervention	Current Status	HS
1	presentation Cre: 0.93 Alb: 2.47 Prot: 1.2	Kidney	46	ADA Anakinra IFX + MTX	Cre: 0.98 Prot: 0.5 Alb: 3.7 FMF and arthritis in remission	Partial Remission
2	Cre: 5.69 (predialysis) Alb: 2.33	Bone marrow	13	None (ADA continued)	ESRD Alb: 2.67	Complete Remission
3	Cre: 0.85 Alb: 2.8 Prot: 7.2	Kidney	24	ACE inhibitor Colchicine (ADA continued)	Cre: 4.85	Partial remission
4	Cre: 4.12-6.78 Alb: 2.4 Prot: 1.6	Gastric	9	IFX	HD discontinued Alb: 2.7 Cre: 3.8 Bladder cancer	Complete Remission
5	Cre: 1,35 Prot:17 Alb: 2.4	Kidney	60	IFX	Cre: 1.42 Alb: 4.07 Pro: 3	Complete Remission
6	Cre: 5.79 Prot: 8 Alb: 1.9	Rectum	13	ADA	ESRD Alb: 2.5	Partial remission
7	Cre: 0.65 Prot: 12 Alb: 2.43	Kidney	8	ADA	Cre: 0.62 Prot: 2.7 Alb: 3.3	Partial remission
8	Cre: 0.79 Prot: 6 Alb: 3.2	Kidney	4	ADA	Cre: 0.71 Prot: 0.8 Alb: 3.8	Complete Remission

Cre: creatinine, Alb: albumin, Prot: 24-hour proteinuria in grams, HD: hemodialysis, ADA: adalimumab, IFX: infliximab, MTX: methotrexate, CsA: cyclosporine, FMF: Familial Mediterranean Fever, ESRD: end-stage renal disease.