	Patients	
	n =90	
Age (years)		
≤39́	19 (21.11)	
40-49	12 (13.33)	
50-59	22 (24.44)	
60-69	14 (15.55)	
70-79	13 (14.44)	
$\geq 80$	10 (11.11)	
Sex		
Male	53 (58.9)	
Female	37 (41.1)	
Smoking		
Yes	11 (12.22)	
No	66 (73.33)	
Previously	11 (12.22)	
Electronic	2 (2.22)	
Allergic rhinitis	7 (7.77)	
Chronic rhinosinusitis	3 (3.33)	
Comorbidities	20 (42 22)	
No medical history	38 (42.22)	
Hypertension	25 (27.77)	
Diabetes	12 (13.33)	
Dyslipidemia	3 (3.33)	
Obesity	3 (3.33)	
Hypothyroidism	3 (3.33)	
Olfactory dysfunction	35 (38.88)	
Gustatory dysfunction	33 (36.66)	
Olfactory and gustatory	20 (22 22)	
dysfunction	50 (55.55)	
Onset of hyposmia		
Before other symptoms	15 (42.86)	
After other symptoms	11 (31.43)	
Nasal blockage	8 (8.89)	
Rhinorrhea	6 (6.66)	
Covid-19 severity		
Mild-Moderate	45 (50)	
Severe	35 (38.88)	
ICU-treated	10 (11.11)	

 Table 1: Demographic and clinical characteristics of the 90

 patients hospitalized for confirmed COVID-19.

Values are given as number of patients and percentage in brackets, ICU: intensive care unit