

**Table 1:** Baseline characteristics of the sixty patients undergoing breast cancer surgery, who were included in this prospective, randomized study, and received serratus plane block alone (Group S) or a combination of pectoral type 1 and serratus plane block (Group SPECS).

	Group S (n =30)	Group SPECS (n =30)	p-value
Age (years)	58.2 (11.6)	54.7 (13.1)	0.292
BMI (kg/m <sup>2</sup> )	29.1 (4.6)	29.9 (4.9)	0.549
Duration of Anesthesia (min)	146.1 (27.0)	148.6 (40.4)	0.785
Duration of Surgery (min)	120.4 (26.4)	120.8 (41.3)	0.962
ASA score			
1	14 (46.7)	10 (33.3)	
2	15 (50.0)	20 (66.7)	0.304
3	1 (3.3)	0 (0.0)	
Type of surgery			
Mastectomy + Axillary dissection	4 (13.3)	6 (20.0)	
Mastectomy + Sentinel Lymph Node Biopsy	20 (66.7)	18 (60.0)	0.777
Modified Radical Mastectomy	6 (20.0)	6 (20.0)	

Values are presented as mean with standard deviation in brackets or count with percentage in brackets, n: number, ASA: American Society of Anesthesiologists physical status, BMI: body mass index.