

**Table 2:** Descriptive values for postoperative nausea-vomiting scale (PONV) score and patients' satisfaction results according to group.

		Group S		Group SPECS		p-value
		n	%	n	%	
PONV 0 (h)	0	10	33.3	23	76.7	<b>0.003</b>
	1	16	53.3	5	16.7	
	2	4	13.3	2	6.7	
PONV 1 (h)	0	10	33.3	21	70.0	<b>0.017</b>
	1	17	56.7	8	26.7	
	2	3	10.0	1	3.3	
PONV 2 (h)	0	15	50.0	26	86.7	<b>0.008</b>
	1	13	43.3	4	13.3	
	2	2	6.7	0	0.0	
PONV 6 (h)	0	24	80.0	27	90.0	0.278
	1	6	20.0	3	10.0	
PONV 12 (h)	0	24	80.0	30	100.0	<b>0.010</b>
	1	6	20.0	0	0.0	
PONV 24 (h)	0	27	90.0	30	100.0	0.076
	1	3	10.0	0	0.0	
Patient Satisfaction	very bad	2	6.7	0	0.0	<b>0.001</b>
	mediocre	18	60.0	2	6.7	
	good	7	23.3	5	16.7	
	very good	3	10.0	23	76.7	

Group S: patients received serratus plane block alone, Group SPECS: patients received a combination of pectoral type 1 and serratus plane block, n: number, PONV: postoperative nausea and vomiting scale (0 to 3 scale).