Table 3: Correlation of the scores of the Covid-19-related questions (range 0-10), of the Hospital Anxiety Depression Scale (HADS; range 0-42), and HADS-Anxiety (HADS-A; range 0-21), HADS-Depression (HADS-D; range 0-21) with the employment status of the study's sample.

Questions/Tests	mean ± SD		
	Employed patients	Unemployed patients	p-value
Fear of getting infected	5.2 ± 2.6	6.6 ± 2.7	0.025*
Fear of death	3.5 ± 3.3	5.6 ± 3.1	0.013*

Fear of getting infected	5.2 ± 2.6	6.6 ± 2.7	0.025*
Fear of death	3.5 ± 3.3	5.6 ± 3.1	0.013*
Fear of sickness for their relatives	7.2 ± 2.1	6.9 ± 3.1	0.077
Stress during lockdown	5.1 ± 3.3	7.3 ± 2.5	0.005*
Fear of AFD unavailability	22 ± 26	3.6 ± 3.1	0.066

Fear of sickness for their relatives	7.2 ± 2.1	6.9 ± 3.1	0.077
Stress during lockdown	5.1 ± 3.3	7.3 ± 2.5	0.005*
Fear of AED unavailability	2.2 ± 2.6	3.6 ± 3.1	0.066
HADS Total	11.4 ± 5.9	18.1 ± 7.1	<0.001*
HADS-A	6.4 ± 3.8	9.4 ± 4.3	0.005*

HADS-D 4.9 ± 2.9 8.7 ± 3.5 < 0.001*

SD: Standard Deviation, AED: Antiepileptic drugs, HADS: Hospital Anxiety Depression Scale, HADS-A: HADS-Anxiety, HADS-D: HADS-

*Values of p in bold are statistically significant (p < 0.05).

Depression.

Unemployed patients scored statistically higher than employed patients in questions regarding fear of getting infected, fear of death, stress during lockdown, HADS total, HADS-A and HADS-D.