

**Table 4 :** Correlation of the scores of Covid-19-related questions (range 0-10), of the Hospital Anxiety Depression Scale (HADS; range 0-42), and HADS-Anxiety (HADS-A; range 0-21), HADS-Depression (HADS-D; range 0-21) with the seizure control status of the study's sample.

Questions/Tests	mean $\pm$ SD		p-value
	seizure-free patients	no seizure-free patients	
Fear of getting infected	5.6 $\pm$ 2.5	6.4 $\pm$ 2.9	0.144
Fear of death	3.8 $\pm$ 3.3	5.6 $\pm$ 3.2	<b>0.034*</b>
Fear of sickness for their relatives	7.2 $\pm$ 2.7	6.9 $\pm$ 2.8	0.761
Stress during lockdown	6.2 $\pm$ 3.3	6.5 $\pm$ 2.8	0.949
Fear of AED unavailability	2.1 $\pm$ 2.5	3.9 $\pm$ 3.1	<b>0.017*</b>
HADS Total	13.3 $\pm$ 6.7	17.1 $\pm$ 7.5	<b>0.026*</b>
HADS-A	7.2 $\pm$ 3.9	9.0 $\pm$ 4.6	0.099
HADS-D	6.1 $\pm$ 3.4	8.1 $\pm$ 3.8	<b>0.025*</b>

\*Values of p in bold are statistically significant ( $p < 0.05$ ).

Patients with incomplete seizure control scored statistically higher than seizure-free patients in questions regarding fear of death, fear of AED unavailability, HADS total, and HADS-D.

SD: Standard Deviation, AED: Antiepileptic drugs, HADS: Hospital Anxiety Depression Scale, HADS-A: HADS-Anxiety, HADS-D: HADS-Depression.