

**Table 1:** Baseline characteristics of the study’s sample consisting of 90 nurses working in Greek intensive care units who participated in the cross-sectional non-interventional study investigating the frequency of various nursing errors and their association with “nurse-related” and “ICU-related” features.

Variable	Study participants
Age (y)	41.6 ± 7.2
Gender (n, %)	
-male	16 (17.8%)
-female	74 (82.2%)
Marital status	
-single	20 (22.2)
-married/with partner	64 (71.1)
-divorced/separated	6 (6.7)
Number of underaged children	2 (0-3)
Working experience in ICU (y)	12.7 ± 7.7
Level of training satisfaction	2.2 ± 0.8
Number of nurses/ICU	29 ± 8.6
Number of beds/ICU	10.8 ± 2.2
Morning shifts/week	2.7 ± 1.2
Evening shifts/week	2.3 ± 1
Night shifts/week	1.9 ± 0.6
Days off work/month	7 ± 1.3
AIS score	6 (0-14)
State anxiety score	40.1 ± 12.5
Trait anxiety score	44.1 ± 9.3
High EE (n, %)	55 (61.1)
High DEP (n, %)	26 (28.9)
Low PA (n, %)	29 (32.2)
High burnout (n, %)	75 (83.3)

Values are presented as means with standard deviation or number with percentage in brackets (where indicated). ICU: Intensive Care Unit, AIS: Athens Insomnia Scale, EE: Emotional Exhaustion, DEP: Depersonalization, PA: Personal Accomplishment.