

**Table 3:** Correlations between demographic and clinical variables regarding study's 58 psoriasis patients.

	<b>Age (years)</b>	<b>Education (years)</b>	<b>Duration of psoriasis (years)</b>	<b>BDI-II</b>	<b>RS</b>	<b>PASI</b>	<b>PGA</b>	<b>BSA</b>
<b>Age (years)</b>	-	-0.314	0.263	0.254	-0.068	-0.151	-0.117	-0.190
		0.016	0.047	0.054	0.614	0.258	0.381	0.154
<b>Education (years)</b>		-	-0.028	-0.114	0.173	0.282	0.243	0.264
			0.837	0.392	0.194	0.032	0.066	0.045
<b>Duration of psoriasis (years)</b>			-	0.034	0.042	0.036	-0.007	0.064
				0.799	0.754	0.787	0.959	0.632
<b>BDI-II</b>				-	-0.517*	-0.042	-0.047	-0.223
					<0.001	0.756	0.729	0.092
<b>RS</b>					-	-0.090	-0.109	0.001
						0.501	0.413	0.992
<b>PASI</b>						-	0.981*	0.941*
							<0.001	<0.001
<b>PGA</b>							-	0.930*
								<0.001
<b>BSA</b>								-

Spearman rho sig. 2-tailed; \*: Statistically significant after Bonferroni corrections for multiple comparisons ( $p < 0.00625$ ). BDI-II: Beck Depression Inventory-II, RS: Resilience Scale, PASI: Psoriasis Area and Severity Index, PGA: Physician Global Assessment, BSA: Body Surface Area.