

Table 3: Cervical conization procedures that the 83 women with cervical pathology underwent.

Histology following conization	n	Number of passes		Depth of cone		Endo-cervical margin *	n
		1	n	≤1cm	n		
LGSIL	1 (1.2)	1	0	≤1cm	0	-	1 (1.2)
		>1	1 (1.2)	>1cm	1 (1.2)	+	0
HGSIL	70 (84.4)	1	47 (56.7)	≤1cm	17 (20.5)	-	50 (60.3)
		>1	23 (27.7)	>1cm	53 (63.9)	+	20 (24.1)
IA1	5 (6)	1	4 (4.8)	≤1cm	1 (1.2)	-	4 (4.8)
		>1	1 (1.2)	>1cm	4 (4.8)	+	1 (1.2)
IA2	2 (2.4)	1	1 (1.2)	≤1cm	0	-	1 (1.2)
		>1	1 (1.2)	>1cm	2 (2.4)	+	1 (1.2)
NEG	5 (6)	1	4 (4.8)	≤1cm	1 (1.2)	-	5 (6)
		>1	1 (1.2)	>1cm	4 (4.8)	+	0
Overall	83 (100)	1	56 (67.5)	≤1cm	19 (23)	-	61 (73.5)
		>1	27 (32.5)	>1cm	64 (77)	+	22 (26.5)

Values are presented as frequencies with percentage in brackets. n: number, LGSIL: low grade squamous intraepithelial lesion, HGSIL: high grade squamous intraepithelial lesion, CIN: cervical intraepithelial neoplasia, IA1: micro-invasive cervical cancer, -: free endocervical margins, +: involved endocervical margins.