

Table 2: Clinical data obtained through the European Cancer Treatment and Organization Committee Quality of Life Questionnaire and the Hospital Anxiety and Depression scale on the first day of radiotherapy and 6 months after the end of radiotherapy.

EORTC QLQ-C30		
Physical Function	first day	66.93 ± 30.05
	6 months	60.26 ± 23.62
Role Function	first day	76.33 ± 28.76
	6 months	64.31 ± 23.06
Emotional Function	first day	67.15 ± 26.64
	6 months	64.12 ± 23.03
Cognitive Function	first day	79.7 ± 21.42
	6 months	72.29 ± 15.78
Social function	first day	80.06 ± 27.08
	6 months	76.2 ± 21.8
Fatigue	first day	33.15 ± 30.91
	6 months	38.3 ± 24.34
Nausea Vomiting	first day	19.93 ± 26.04
	6 months	17.71 ± 20.77
Pain	first day	23.53 ± 29.09
	6 months	30.46 ± 27.68
Dyspnea	first day	41.22 ± 12.3
	6 months	40.92 ± 27.78
Insomnia	first day	37.26 ± 36.29
	6 months	29.8 ± 31.19
Appetite Loss	first day	28.10 ± 35.51
	6 months	30.46 ± 29.43
Constipation	first day	23.52 ± 32.17
	6 months	38.3 ± 33.41
Diarrhea	first day	18.31 ± 29.29
	6 months	25.23 ± 25.09
Financial difficulty	first day	18.95 ± 32.15
	6 months	35.69 ± 38.04
Global Health	first day	40.69 ± 28.41
	6 months	30.78 ± 16.02
HADS scale		
Depression	first day	10.22 ± 3.62
	6 months	9.67 ± 3.63
Anxiety	first day	8.33 ± 3.06
	6 months	8.96 ± 2.77

Values are presented as mean ± standard deviation. EORTC QLQ-C30: European Cancer Treatment and Organization Committee Quality of Life Questionnaire, HADS: Hospital Anxiety and Depression Scale.